

# THE HOMOTOPY TYPE OF THE COMPLEMENT OF THE CODIMENSION-TWO COORDINATE SUBSPACE ARRANGEMENT

JELENA GRBIĆ AND STEPHEN THERIAULT

A complex coordinate subspace of  $\mathbb{C}^n$  is given by

$$L_\sigma = \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid z_{i_1} = \dots = z_{i_k} = 0\}$$

where  $\sigma = \{i_1, \dots, i_k\}$  is a subset of  $[n]$ . For each simplicial complex  $K$  on the set  $[n]$  we associate the complex coordinate subspace arrangement  $\mathcal{CA}(K) = \{L_\sigma \mid \sigma \notin K\}$  and its complement  $U(K) = \mathbb{C}^n \setminus \bigcup_{\sigma \notin K} L_\sigma$ . On the other hand, to  $K$  we can associate the Davis-Januszkiewicz space  $DJ(K) = \bigcup_{\sigma \in K} BT_\sigma \subset BT^n$ , where  $BT^n$  is the classifying space of  $n$ -dimensional torus, that is, the product of  $n$  copies of infinite-dimensional projective space  $\mathbb{C}P^\infty$ , and  $BT_\sigma := \{(x_1, \dots, x_n) \in BT^n \mid x_i = * \text{ where } i \notin \sigma\}$ . Let  $\mathcal{Z}_K$  be the fibre of  $DJ(K) \rightarrow BT^n$ . By [BP, 8.9], there is an equivariant deformation retraction  $U(K) \rightarrow \mathcal{Z}_K$ , and the integral cohomology of  $\mathcal{Z}_K$  has been calculated in [BP, 7.6 and 7.7].

**Theorem 1.** *The complement of the codimension-two coordinate subspace arrangement in  $\mathbb{C}^n$  has the homotopy type of the wedge of spheres*

$$\bigvee_{k=2}^n (k-1) \binom{n}{k} S^{k+1}.$$

*Proof.* Let  $K$  be a disjoint union of  $n$  vertices. Then  $DJ(K)$  is the wedge of  $n$  copies of  $\mathbb{C}P^\infty$  and  $U(K)$  is the complement of the set of all codimension-two coordinates subspaces  $z_i = z_j = 0$  for  $1 \leq i < j \leq n$  in  $\mathbb{C}^n$ . Therefore to prove the theorem we have to determine the homotopy fibre of the inclusion  $\bigvee_{t=1}^n \mathbb{C}P^\infty \rightarrow \prod_{t=1}^n \mathbb{C}P^\infty$ . This is done by applying Proposition 5 to the case  $X_1 = \dots = X_n = \mathbb{C}P^\infty$  and noting that  $\Omega \mathbb{C}P^\infty \simeq S^1$ .  $\square$

It should be emphasized that Theorem 1 holds without suspending. Previously, decompositions were known only after some number of suspensions, the best of which was by Schaper [S] who required one suspension. To finish the proof of Theorem 1 it remains to prove Proposition 5. This was originally proved by Porter [P] by examining subspaces of contractible spaces. We present an accelerated proof based on the Cube Lemma.

We work in the category of based, connected topological spaces and continuous maps. Let  $*$  denote the basepoint. For spaces  $X, Y$ , let  $X \rtimes Y = (X \times Y)/(* \times Y)$ ,  $X \wedge Y = (X \rtimes Y)/(X \times *)$ , and  $X * Y = \Sigma X \wedge Y$ . Denote the identity map on  $X$  by  $X$ . Denote the map which sends all points to the basepoint by  $*$ .

**Lemma 2.** *Let  $A, B$ , and  $C$  be spaces. Define  $Q$  as the homotopy pushout of the map  $A \times B \xrightarrow{* \times B} C \times B$  and the projection  $A \times B \xrightarrow{\pi_1} A$ . Then  $Q \simeq (A * B) \vee (C \rtimes B)$ .*

*Proof.* Consider the diagram of iterated homotopy pushouts

$$\begin{array}{ccccc} A \times B & \xrightarrow{\pi_2} & B & \xrightarrow{i_2} & C \times B \\ \downarrow \pi_1 & & \downarrow * & & \downarrow s \\ A & \xrightarrow{*} & A * B & \xrightarrow{t} & \overline{Q} \end{array}$$

where  $\pi_2, i_2$  are the projection and inclusion respectively. Here, it is well known that the left square is a homotopy pushout, and the right homotopy pushout defines  $\overline{Q}$ . Note that  $i_2 \circ \pi_2 \simeq * \times B$ . The outer rectangle in an iterated homotopy pushout diagram is itself a homotopy pushout, so  $\overline{Q} \simeq Q$ . The right pushout then shows that the homotopy cofibre of  $C \times B \rightarrow Q$  is  $\Sigma B \vee (A * B)$ . Thus  $t$  has a left homotopy inverse. Further,  $s \circ i_2 \simeq *$  so pinching out  $B$  in the right pushout gives a homotopy cofibration  $C \rtimes B \rightarrow Q \xrightarrow{r} A * B$  with  $r \circ t$  homotopic to the identity map.  $\square$

**Lemma 3.** *Let  $Y_1, \dots, Y_n$  be spaces. Then there is a homotopy equivalence*

$$\Sigma(Y_1 \times \dots \times Y_n) \simeq \bigvee_{k=1}^n \left( \bigvee_{1 \leq i_1 < \dots < i_k \leq n} \Sigma Y_{i_1} \wedge \dots \wedge Y_{i_k} \right).$$

*Proof.* Induct on the decomposition  $\Sigma(A \times B) = \Sigma A \vee \Sigma B \vee (\Sigma A \wedge B)$ . □

The following was proved by Mather [M] and is known as the Cube Lemma.

**Lemma 4.** *Suppose there is a diagram of spaces and maps*

$$\begin{array}{ccccc} E & \xrightarrow{\quad} & F & & \\ & \searrow & \downarrow & \searrow & \\ & & G & \xrightarrow{\quad} & H \\ & \downarrow & \downarrow & \downarrow & \downarrow \\ A & \xrightarrow{\quad} & B & & \\ & \searrow & \downarrow & \searrow & \\ & & C & \xrightarrow{\quad} & D \end{array}$$

where the bottom face is a homotopy pushout and the four sides are obtained by pulling back with  $H \rightarrow D$ . Then the top face is a homotopy pushout. □

**Proposition 5.** *Let  $X_1, \dots, X_n$  be spaces. Consider the homotopy fibration*

$$F_n \rightarrow X_1 \vee \dots \vee X_n \rightarrow X_1 \times \dots \times X_n$$

obtained by including the wedge into the product. Then there is a homotopy decomposition

$$F_n \simeq \bigvee_{k=2}^n \left( \bigvee_{1 \leq i_1 < \dots < i_k \leq n} (k-1)(\Sigma \Omega X_{i_1} \wedge \dots \wedge \Omega X_{i_k}) \right).$$

*Proof.* We induct on  $n$ . When  $n = 2$  it is well known that  $F_2 \simeq \Sigma \Omega X_1 \wedge \Omega X_2$ . Let  $n \geq 3$  and assume the Proposition holds for  $F_{n-1}$ . Let  $M_k = X_1 \vee \dots \vee X_k$  and  $N_k = X_1 \times \dots \times X_k$ . Observe that  $M_n$  is the pushout of  $M_{n-1}$  and  $X_n$  over a point. Composing each vertex of the pushout into  $N_n$  we obtain homotopy fibrations  $\Omega N_n \rightarrow * \rightarrow N_n$ ,  $\Omega N_{n-1} \rightarrow X_n \rightarrow N_n$ ,  $F_{n-1} \times \Omega X_n \rightarrow M_{n-1} \rightarrow N_n$ , and  $F_n \rightarrow M_n \rightarrow N_n$ . Write  $N_n$  as  $N_{n-1} \times X_n$ . Then Lemma 4 implies that there is a homotopy pushout

$$\begin{array}{ccc} \Omega N_{n-1} \times \Omega X_n & \xrightarrow{h} & F_{n-1} \times \Omega X_n \\ \downarrow g & & \downarrow \\ \Omega N_{n-1} & \xrightarrow{\quad} & F_n \end{array}$$

where  $g$  is easily identified as the projection and  $h$  is the connecting map for the homotopy fibration  $F_{n-1} \times \Omega X_n \rightarrow M_{n-1} \times * \rightarrow N_{n-1} \times X_n$ . So  $h \simeq \partial_{n-1} \times \Omega X_n$  where  $\partial_{n-1}$  is the connecting map of the fibration  $F_{n-1} \rightarrow M_{n-1} \rightarrow N_{n-1}$ . But  $\partial_{n-1} \simeq *$  as  $\Omega M_{n-1} \rightarrow \Omega N_{n-1}$  has a right homotopy inverse. Thus  $h \simeq * \times \Omega X_n$ . By Lemma 2,  $F_n \simeq (\Omega N_{n-1} * \Omega X_n) \vee (F_{n-1} \times \Omega X_n)$ . Since  $F_{n-1}$  is a suspension,  $F_{n-1} \times \Omega X_n \simeq F_{n-1} \vee (F_{n-1} \wedge \Omega X_n)$ . Combining the decomposition of  $\Sigma \Omega N_n \simeq \Sigma(\Omega X_1 \times \dots \times \Omega X_n)$  in Lemma 3 with the inductive decomposition of  $F_{n-1}$  and collecting like terms, the asserted wedge decomposition of  $F_n$  follows. □

The authors wish to thank Taras Panov and Nigel Ray for encouraging us to consider this problem.

## REFERENCES

- [BP] V.M. Buchstaber and T.E. Panov, *Torus actions and their applications in topology and combinatorics*, University Lecture Series **24**, American Mathematical Society, (2002).
- [M] M. Mather, Pull-backs in homotopy theory, *Canad. J. Math.* **28** (1976), 225-263.
- [P] T. Porter, The homotopy groups of wedges of suspensions, *Amer. J. Math.* **88** (1966), 655-663.
- [S] Ch. Schaper, Suspensions of affine arrangements, *Math. Ann.* **309** (1997), 463-473.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ABERDEEN, ABERDEEN AB24 3UE, UNITED KINGDOM  
*E-mail address:* jelena@maths.abdn.ac.uk  
*E-mail address:* s.theriault@maths.abdn.ac.uk